



Effect of Different Levels of Poultry Manure on Growth and Yield of a Maranthus Plant (*Amaranthus hybridus*) in Obio Akpa, Akwa Ibom State, Nigeria

Akata, O. R ^{a*}, Essien, O. A ^b, Ben, F. E ^a, Enyong, J. K. ^a
and Tochukwu, N. E ^a

^a Department of Crop Science, Akwa Ibom State University, Obio Akpa Campus, Akwa Ibom State, Nigeria.

^b Department of Soil Sciences, Akwa Ibom State University, Obio Akpa Campus, Akwa Ibom State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajsspn/2024/v10i4445>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/128367>

Original Research Article

Received: 14/10/2024

Accepted: 18/12/2024

Published: 27/12/2024

ABSTRACT

The field experiment was carried out at the Teaching and Research farm of Akwa Ibom State University Obio Akpa campus, in 201 B to determine the Optimum Poultry manure rate for Amaranthus. The experiment was laid out in a randomized complete block design replicated three

*Corresponding author: E-mail: akatafonime@aksu.edu.ng;

Cite as: O. R, Akata, Essien, O. A, Ben, F. E, Enyong, J. K., and Tochukwu, N. E. 2024. "Effect of Different Levels of Poultry Manure on Growth and Yield of a Maranthus Plant (*Amaranthus Hybridus*) in Obio Akpa, Akwa Ibom State, Nigeria". *Asian Journal of Soil Science and Plant Nutrition* 10 (4):744-51. <https://doi.org/10.9734/ajsspn/2024/v10i4445>.

times. The treatments were four rates of poultry manure and 6 t/ha. Growth and yield parameters were assessed, and data collected were subjected to analysis of variance (ANOVA). Significance difference at 5% probability. Results showed that poultry manure rate had significant (p < 0.05) effects on all the growth and yield parameters assessed except stem girth. The treatment that received 6 t/ha performed best in all the parameters, while the least was recorded from the control treatment. The application of 6 t/ha poultry manure, produced 12.19 t/ha fresh foliage yield, followed by the plot that received 4 t/ha poultry (11.33 t/ha), The treatment of 2 t/ha produced 7.70 t/ha foliage yield. The lowest foliage yield of 3.18 t/ha was harvested from the control plot the application of poultry manure also enhanced dry matter accumulation. The plot of 6 t/ha poultry manure produced 30.71g matter, while the control treatment had 33.14g. Therefore, the study recommended the application of 4t/ha poultry manure for optimum fresh foliage yield and dry matter accumulation.

Keywords: Poultry manure; different rate; growth yield parameter; dry matter; harvesting interval; Amaranthus plant.

1. INTRODUCTION

Amaranthus is a perennial vegetable. Amaranthus is a cosmopolitan genus of herbs. The genus Amaranthus, in the family Amaranthaceae has more than 800 species. These include *Amaranthus hybridus*, *Amaranthus cruetus*, and others are consumed as leaf vegetables [1]. Amaranthus originated in America and is one of the oldest food crops in the world, with evidence of its cultivation reaching back as far as 6700 BC. At present, Amaranthus is extensively grown as a green, leafy vegetable in many temperate and tropical regions [2]. The largest area ever grown was during the height of the Aztec civilization in Mexico during the 1400s. After the arrival of the Spanish conquistadors in Mexico in the early 1500s, Amaranthus almost disappeared from America as a crop until research began on it in the US in the 1970s. Amaranthus has spread around the world (Including Mexico, Central America, India, and Nepal).

According to Akubugwo [3], approximate composition Of Amaranthus hybrids L. Leaves is moisture content 83.48%, Ash, Content 1.380%, crude protein 17.92%, crude oil 4.65%, crude fiber 8.61%, available carbohydrate 52.18 and calorific value (Kcal/ 100g) 268.92 [4]. Nigeria has also been claimed as the center of diversity, corroborated by the prevalent use of local names and the enormous genetic diversity available there. Amaranthus plays important roles in nutrition, it provides calcium, iron, Vitamin A (carotene) Vitamin C (ascorbic acid), [5]. It is used as a refrigerant, diuretic, and purgative, as an enema for stomach troubles, and piles, against cholera, and also as a sudorific. People with hypertension and

cardiovascular diseases can use the leaves as an effective alternative to drug therapy [3,6]. Their nutrition quality receives world attention, especially as a rich source of minerals and protein [7,8].

Some Amaranthus species are also cultivated for their seeds, which are known to contain about 16% protein and are consequently utilized in confectionaries. Approximately 60 species are presently recognized with inflorescence and foliage ranging from purple and red to gold. The leaf vegetable has a rich profile of essential body metabolites such as vitamins A, B, and C mineral elements such as iron, calcium, and phosphorus, and some measure of protein [9,10]. It is principally grown for its large fleshy leaves and herbaceous stem which are succulent palatable and nutritious. It can be boiled and added to soups or stews and is also useful for livestock feed and human consumption. Invariably, the leaves of grain amaranthus could be used for feeding rabbits and other micro livestock at the early stages of growth [5,11].

Amresh et al. [12] reported hepatoprotective and antioxidant activity of 50% ethanol extract of the whole plant of Amaranthus spinosus against carbon tetrachloride (CC4) induced. The crop is adapted to a wide range of soil conditions. Sandy soil with slightly acidity is best suited. A temperature range of 20-30°C is required, the land should be plowed or dug followed by leveling then shallow trenches of width 30-35cm, well rotten farmyard manure is mixed with soil in the trenches [13]. Amaranthus is grown during both wet and dry seasons though irrigation is normally required for dry season crops, since the rate of transpiration by the leaves is fairly

high. It prefers soil with a pH above 6. Amaranths are planted either by direct seeding or transplanting. Since amaranths seeds are very small, it is mixed with sand at a ratio of 1g seed and obtain a uniform Stand, then cover seed highly with a layer of compost or rice hulls immediately after broadcast [14,15].

In crop production, organic and inorganic materials such as poultry manure are used to supplement chemical fertilizers [16,17,18]. Poultry manure like other manure is known to improve the soil by enhancing both physical and chemical properties. According to Okoli et al. [19] and Ekwere et al. [20], the use of poultry manure in amaranthus production is desirable as it has variable impacts on most parameters assessed, the result of the study showed that the poultry manure rates are capable of improving of the yield of Amaranthus. Nweke and Obasi [21] observed that higher organic manure rates 15 t/ha and 20 t/ha, and pig manure rates 12 t/ha and 24 t/ha, respectively gave higher yield results than where manure was not applied

According to Grubben [22], Amaranthus species are endowed with a rapid growth rate which could encourage shoot re-growth following cutting. Furthermore, Ben et al. [23] and Essien et al. [24] viewed increasing nutrient content is by boosting with organic material; poultry manure. Hence, there is a need to research the effect of different levels of poultry manure on the growth and yield of the amaranthus plant.

2. MATERIALS AND METHODS

2.1 Experimental Site

This research was carried out between July to October, 2018 at the Teaching and Research Farm, Obio Akpa Campus, Akwa Ibom State University, Nigeria. The research site is situated between latitude 4° 31' 1" and 5° 30' N and longitude 8° 30' and 8° 00' E, with an annual rainfall of 2500mm. The total area of land for the experiment was 15m x 17m (255 m²). Twelve beds were made with the dimensions of 4m X 3m each with pathways of 1m within the land area and 0.05m between other land areas.

2.2 Experimental Treatments

Four treatments were replicated three times which included three different levels of Poultry manure and one in which no poultry manure was added used as control, that is C tons ha⁻¹, 2 tons ha⁻¹, 4 tons ha⁻¹, and 6 tons ha⁻¹. The poultry

manures were obtained from the University poultry farm.

2.3 Nursery Management

Amaranthus black seeds were sown in prepared nursery seedbeds on the experiment site. A small quantity of manure was applied in the seedbeds to aid its germination, seeds were sown by broadcasting seeds over the bed. Transplanting was done in the evening to reduce temperature stress.

2.4 Experiment Design

The experiment was laid out in a randomized Complete Block Design (RCBD) with three replications.

2.5 Cultural Practices

Land preparation: The land was cleared and debris was packed.

Transplanting: After 30 days in the nursery, the seedlings were transplanted. The transplanting was done at a spacing of 50cm x 50cm.

Weeding: This was done twice at 2 weeks after transplanting (WAT) and at 6 weeks after transplanting (WAT).

Fertilizer Application: Poultry manure was applied at 2 weeks and 6 weeks after transplanting using ring method.

Data Collection: Data were collected on the field from four (4) randomly selected plants and were tagged from each plot.

The growth parameters measured were;

Plant Height (cm): The plant height was measured with the aid of a meter rule from the soil level to the plant apex.

Stem Girth (cm): This was measured by tying a thread around the base area and reading the marks of a ruler.

Leaf Area (cm²): This was determined by measuring the length and width of the leaf of tagged plants. It was later corrected factor.

Thus $LA = 0.8 \times LW$ (Kuet and svobada, 1971)

Where

L=Length of the leaf

W=Width of leaf

Number of leaves: This was determined by manual counting of the marketable leaves on the plant.

Number of Branches: The number of branches was carefully counted on each tagged plant. The yield components were:

Dry Matter Accumulation (g): This was determined by removing the moisture content of fresh foliage either through Sun drying or oven drying.

Foliage Weight (t/ha): This can be determined by cutting the leaves of the amaranthus plant and weighed.

Germination Percentage (%): The germination percentage was determined by dividing the unsuccessful transplants from the successful transplants and multiplying the total by the percentage mark (100). The number of unsuccessful plants was out of 576 plants, given a germination percentage of 83.33% was recorded.

2.6 Statistical Analysis

Data collected were subjected to analysis of variance (ANOVA) to determine the level of significance of treatment and means were significantly separated by the use of Least Significant Difference (LSD) at a 5% probability level.

3. RESULTS AND DISCUSSION

3.1 Results

Result of soil physico-chemical properties at the Experimental Site before planting (Table 1), The result showed that the soil was sandy soil. Organic carbon and total N content were low [25,26]. The soil was acidic [25,27] Therefore, the soil growth yield required the application of manure for proper yield.

Number of Amaranthus leaves Per Plant as Influenced by Poultry Manure Rates: The number of leaves per plant of Amaranthus was significantly influenced by poultry manure application at 4, 6, 8, 10, and 12 weeks after

transplanting (WAT), shown in Table 2. The amaranthus that received the higher dose of poultry manure (4 and 6 t/ha) was significantly taller than those that received 2t/ha manure, which were also taller than the control. The treatment of 6 t/ha had 38.40, 70.33, 97.52, 103.42, and 119.55 leaves per plant at 4,6,8,10 and 12 WAT, respectively. The control treatment had 14.60, 27.40, 33.70, 38.30 and 46.40 leaves per plant, respectively.

Table 1. Soil Physico- chemical properties of the experimental site before planting

Soil Properties	
Sand	88.70
Silt	6.30
Clay	5.00
PH (H ₂ O)	5.10
Electrical conductivity (ds/m)	0.08
Total Nitrogen (%)	0.11
Available phosphorus	87.55
Organic Carbon (%)	1.42
Exchangeable Ca	3.01
Exchangeable Mg	1.59g
Exchangeable K	0.11
Exchangeable Na	0.07
Exchangeable Acidity	1.95
ECEC	6.73
Base Saturation (%)	71.03

ECEC- Effective Cation Exchangeable Capacity

Amaranthus Height (cm) as Influenced by Poultry Manure Rates: Plant height as an influence by poultry manure rates indicated a significant difference in all the weeks under investigation except at 2 WAT- The increase in poultry manure rate from 2 to 4 t/ha indicated a significant increase in plant height (Table 3), whereas statistically there was no significant difference observed between the plant height in 4 t/ha and 6 t/ha rates, at 12 WAT- The plant that received 6 t/ha was significantly different from the height in 2t/ha.

Leaf Area of Amaranthus as Influenced by Poultry Manure Rates: Leaf area as affected by Poultry manure rates varied significantly at 4, 6, 8, 10, and 12 WAT (Table 4). The largest leaf area, 91.33, 100.09, 113.70, 95.01, and 94.83 cm² at 4, 6, 8, 10, and 12 WAT respectively. The leaf area recorded in the

treatment of 6 t, the poultry manure rate was not statistically significant, when compared to 91.30, 98.30, 110.66, 9, 3.75, and g3.70, respectively, recorded from the treatment of 4 t/ha with the leaf area of 35.02, 29.15, 25.60, 20.48 and 20.11 cm, respectively.

Stem Girth of Amaranthus as Influenced by Poultry Manure Rates: Stern girth as affected by poultry manure was not significantly different ($P > 0.05$) in all the sample weeks (Table 5). However, in the stem girth, the highest dose of poultry manure treatment recorded the biggest stem girth and was statistically significant when compared to the mean values recorded in the control treatment.

Number of branches per plant as influenced by poultry manure rates: The number of branches per plant differed significantly ($P > 0.05$) at 4,6,8,10 and 12 WAT (Table 6). The result showed that the Amaranthus plants that received the higher dose of poultry manure (4 and 6t/ha) branched profusely compared to

the number of branches in 2t/ha plots and control treatment.

The result further indicated a significant increase in the number of branches per plant when 2t/ha rate was compared to the control plot. At 12 WAT, the plot that received 6 t/fha poultry manure had 31.75 branches, while 28.60 and 21.77 were recorded in 4 t/ ha and 2t/ha poultry manure rates, respectively. The least number of branches per plant (14.06) was observed from the control treatment.

Total Fresh Foliage Yield and Dry Matter Accumulation: Total foliage yield as influenced by poultry manure rate varied significantly ($P < 0.05$) when compared to 7.70 t/ ha and 3.18 t/ha from 2t/ha poultry manure and control plots respectively. The total fresh foliage yield obtained from the treatment that received 6t/ ha poultry manure (12.19 t/ha) was not significantly different ($+0.05$) when compared to 11.33 t/ha total fresh foliage yield harvested from the treatment of 4 t,fha poultry manure.

Table 2. Number of leaves per plant as influenced by poultry manure

Rates of Poultry Manure (t/ha)	Weeks after Transplanting (WAT)					
	2	4	6	8	10	12
Control	780	14.6	27-40	33.72	38.30	46.40
2	8.33	22.46	51.70	71.50	80.44	88.21
4	9.40	34.60	68.09	91.30	101.01	12.40
6	9.42	38.40	70.33	97.52	103.42	119.55
LSD ($P > 0.05$)	NS	3.71	5.00	5.39	6.50	6.80

* NS Not Significant

Table 3. Amaranthus height as influenced by poultry manure rate

Poultry manure Rate (t/ ha)	Weeks After Transplanting (WAT)					
	2	4	6	8	10	12
Control	16.75	24.46	26.31	29-40	31.01	35.73
2	15.88	46.67	49.25	53.11	53.14	61.06
4	16.42	57.75	52.36	56.91	61.33	63.9B
6	16.45	58.33	52.54	57.20	64.02	64.51
CSO ($p > 0.05$)	NS	2.62	2.70	3.01	3.18	3.25

NS = Not Significant

Table 4. Leaf Area (Cm²) of Amaranthus as influenced by poultry manure rates

Poultry manure Rate (t/ha)	Weeks After Transplanting (WAT)					
	2	4	6	8	10	12
Control	13.93	35.02	29.15	25.60	20.48	20.11
2	13.99	50.85	53.01	39.65	39.64	39.25
4	14.01	91.30	98.32	110.66	93.75	93.70
6	13.97	91.33	100.09	113.70	95.01	94.33
LSD ($P < 0.05$)	NS	2.17	3.70	3.92	3.42	3.20

NS: Not Significant

Table 5. Stem girth of Amaranthus as influenced by poultry manure rates

Poultry manure	Weeks After Transplanting (WAT)					
	2	4	6	8	10	12
Rate (t/ha)	2	4	6	8	10	12
Control	1.71	2.84	3.73	3.78	3.79	3.81
2	1.75	3.71	4.81	4.88	4.89	5.03
4	1.76	3.75	4.98	5.06	5.08	5.10
6	1.73	3.76	5.10	5.11	5.15	5.17
LSD (P<0.05)	NS	NS	NS	NS	NS	NS

NS Not Significant

ID

Table 6. Number of branches per plant as influenced by poultry manure rates

Poultry manure Rate (t/ha)	Weeks After Transplanting (WAT)					
	2	4	6	8	10	12
Control	0.33	2.89	5.75	9.72	12.41	14.06
2	0.33	4.55	8.06	14.98	19.34	21.77
4	0.32	6.33	10.31	19.11	24.67	28.60
6	0.36	6.81	12.41	22.40	26.40	31.75
LSD (P<0.05)	NS	1.60	1.71	2.18	2.36	3.23

NS = Not significant

Table 7. Foliage yield and dry matter accumulation (g) of amaranthus as influenced by poultry manure rates

Poultry manure rate (t/ha)	Total fresh foliage (t/ha)	Dry matter accumulation 1
Control	3.1 B	23.14
2	7.70	29.25
4	11.33	30.71
6	12.19	30.71
LSD (P<0.05)	1.88	1.51

obtained from battery cages. The increase in growth and yield of Amaranthus could be due to higher nutrient content in the manure [30,31].

The increase in vegetative growth with an increase in manure rate could be that the nitrogen content of the poultry manure promotes vegetative growth of the plants, this observation conforms with Akata et al. [29]; Ikeh et al. [32] that N enhanced the photosynthesis process in turn resulted in high vegetative yield [33].

3.2 Discussion

The significant number of leaves per plant, leaf area, plant height, total fresh foliage yield, and dry matter accumulation in the plots with poultry manure rates could be attributed to the improvement of soil fertility which in turn increased the vigour in terms of the plants. The findings of this investigation are in consonance with the report of Ogedegbe [28]; and Akata [29] that an increase in the dose of poultry manure significantly increases the growth and fresh foliage yield of Amaranthus crenatus.

The poultry droppings obtained were from battery cage houses, which contain higher nitrogen than those from deep litter houses, where sawdust and wood shaving constitute a significant part of total manure, thereby having a high CN ratio and a lower rate of mineralization (Akata, 2006).

Ogedegbe [28] and Akata [29] reported higher Amaranthus yield in plots treated with manure

4. CONCLUSION

The application of poultry manure enhanced the growth and yield of Amaranthus. The fresh foliage yield and dry matter accumulation obtained from the treatment of 6t/ha were significantly different when compared to 2t/ha and control treatments. Furthermore, the fresh foliage yield and dry matter accumulation recorded from the treatment that received 6t/ha and 4 t/ha poultry manure were not significantly different. It is recommended that the application of 4t/ha poultry manure should be appropriate for optimum production of Amaranthus in Obio Akpa soils of Akwa Ibom State.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Findall HD. Commercial Vegetable Growing. Oxford Tropical Handbook. Oxford: Oxford University Press; 1975.
- Khan AU. Diseases management in the agriculture plants and crops. In: Insect Pests, Diseases, and Weeds Management. 6376:1–46. Available:<https://easychair.org/publications/preprint/zWpF>
- Akubugwo IE, Obasi NA, Chinyere GC, Igbogu AE. Nutritional and chemical value of *Amaranthus hybridus* L. leaves from Afikpo, Nigeria. Afr J Biotechnol. 2007;6(24):2833–9.
- Khan AU. Insect pests management in the agriculture plants and crops. In: Insect Pests, Diseases, and Weeds Management. 6375:1–50. Available:<https://easychair.org/publications/preprint/bsnk>
- Palada MC, Chang LC. AVRDC International Cooperators Guide. Suggested Cultural Practices for Vegetable Amaranth. Available:<http://www.avrdc.org>
- Khan AU. Weeds management in the agriculture plants and crops. In: Insect Pests, Diseases, and Weeds Management. 6377:1–35. Available:<https://easychair.org/publications/preprint/7xDW>
- Olufolaji AO, Tayo TO. Growth, development, and mineral contents of three cultivars of *Amaranthus*. Sci Hortic. 1980;13:131–9.
- Ikeh AO, Ndaeyo NU, Akpan EM, Lidoh EI, Akata OR. Evaluation of complementary use of organic manure for sustainable water yam production in Oyo, Southeastern Nigeria. Am J Res Commun. 2013;1(2):33–48.
- Omueti O. Effects of age of *Celosia* cultivars. Exp Agric. 1980;16:279–86.
- Esang DM, Akata OR, Ikeh AO. Adaptability of seven maize (*Zea mays* L.) varieties in the high humid rainforest zone of Nigeria. Am J Agric Sci Eng Technol. 2021;5(2):92–101.
- Akpaninyang FE, Opara EU. The influence of toxins in disease symptom initiation in plants: A review. J Agric Sustain. 2017;10(1).
- Amresh G, Zeashan H, Singh S, Rao C. Hepatoprotective and antioxidant activity of *Amaranthus spinosus* against CCl₄-induced toxicity. J Ethnopharmacol. 2009;125:364–6.
- Costea M, Demason D. Stem morphology and anatomy in *Amaranthus* L. (Amaranthaceae). Taxonomic significance. J Torrey Bot Soc. 2007;128:254–61.
- AVRDC. Progress Report 2003. Shanhua, Tainan: Asian Vegetable Research and Development Center; 2003.
- Udounang PI, Ekwere OJ, Akpaninyang FE. Comparative effect of fertilization on cocoyam and component crops in cocoyam-based intercropping systems in Uyo, Southeastern Nigeria. Int J Agric Policy Res. 2022;10(1):16.
- Akpan E, Essien OA, Akata OR, Opeolu EM. Effect of oil palm bush ash and harvesting intervals on okra (*Abelmoschus esculentus* L.) cultivars in Obio Akpa, Akwa Ibom State, Nigeria. Niger J Agric Food Environ. 2024;20(1):21–30.
- Khan AU, Ema IJ, Afsana AS, Khan AU. Home garden and women participation: A mini review. Curr Res Agric Farming. 2021;2(4):46–52. Available:<https://doi.org/10.18782/2582-7146.152>
- Akata OR, Essien OA, Akpan EA, Akpan EG. Effect of inorganic fertilizer and lime rate on chemical properties of soil and yield of cassava (*Manihot esculenta* Crantz) in Obio Akpa, Akwa Ibom State. J Agric Environ Resour Manag. 2024;6(5):48–60.
- Okoli PSO, Nweke IA. Effect of different rates of poultry manure on growth and yield of *Amaranthus* (*Amaranthus cruentus*). IOSR J Agric Vet Sci. Available from: www.iosrjournals.org
- Ekwere OJ, Ilmoh FO, Essien OA. Effect of poultry manure and lime on growth and yield of okra. Mgbenan J Crop Sci. 2022;8(1):15–22.
- Nweke IA, Obasi MN. Effect of different levels of pig manure on the growth and yield of okra. Proc 47th Annu Conf Agric Soc Niger, Ibadan; 2013. p. 23–6.
- Grubben GJH. Cultivation of *Amaranthus* as a tropical leaf vegetable with special reference to southern Dahomey. Commun Dep Agric Res Royal Trop Inst. 1976;67:65–80.

23. Ben FE, Okpara DA, Akpan EA, Udounang PI, Akata OR, Inyang P. Response of orange-fleshed sweetpotato (*Ipomoea batatas* (L.) Lam) to ratoon cropping and nutrient management in Southern Nigeria. *Int J Agric Policy Res.* 2024;12(2):23.
24. Essien OA, Akpan EA, Akata OR. Comparative methods of land use practices effect on clay dispersion in coastal plain sand soils in Southeastern Nigeria. *J Agric Environ Resour Manag.* 2023;5(6):1022–320.
25. Essien OA, Inyang EA, Simeon SO. Aggregate-size distribution: Impact on bulk density and hydraulic conductivity of soil derived from two parent materials in South South Nigeria. *Niger J Agric Food Environ.* 2024;20(1):78–89.
26. Umoh FO, Essien OA, Osodeke VE, Amadi DJ. Evaluation of phosphorus fixation capacity on river alluvium, beach ridge, and coastal plain sand of Akwa Ibom State, Nigeria. *J Agric Ecosyst Manag.* 2021a;1(2):23–8.
27. Umoh FO, Essien OA, Ijah CJ, Sunday AE. Determination of phosphorus fixation capacity on alluvium, sandstone, and shale soils of Akwa Ibom State, Nigeria. *Int J Sustain Agric Res.* 2021b;8(2):105–13.
28. Ogedegbe SA, Ogah JJ, Ajaja BA. Response of *Amaranthus* spp. varieties to organic fertilizer in Vom, Nigeria. University of Nigeria, Nsukka; 2013. p. 90–6.
29. Akata OR, Uko AE, Nwagwu FA, Ndaeyo NU, Ikeh AO, Essang DM. Weed dynamics, growth, and yield of cassava (*Manihot esculenta* Crantz) as influenced by organic fertilizer sources and rates in an Ultisol. *J Basic Appl Res Int.* 2016;18(2):68–76.
30. Akpaninyang F, Okpara DA, Njoku JC, Ogbologwung P, Okpara B, Akpan AE, Jan L. Combinations of organic and inorganic fertilizers for vine, carotene, and root yields of orange-fleshed sweetpotato at Umudike, southeastern Nigeria. *Niger Agric J.* 2015;46(1):87–94.
31. Ekwere OJ, Akpan EA, Akata OR. Performance and yield of fluted pumpkin (*Telfairia occidentalis* (f.) Hook) on an Ultisol amended with soybean meal. *J Agric Environ Resour Manag.* 2019.
32. Ikeh AO, Orji JO, Sampson JO, Akata OR. Effect of oil palm bunch refuse ash in sustainable production of Egusi-melon (*Colocynthis citrullus*) in an Ultisol. *Asian J Agric Hort Res.* 2023;10(3):138–48.
33. Akpan EA, Akpan AA, Elijah, Essien OA. Response of carrot (*Daucus carota* L.) to different organic fertilizers in Obio Akpa, Akwa Ibom State. *J Agric Technol.* 2021;13:46–52.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/128367>